

# Consider ErecAid® for Patients who Fail PDE5i Monotherapy



Millions of men have erectile dysfunction and try PDE5-inhibitor therapy<sup>1</sup>

Not all patients achieve desired results with PDE5 inhibitors<sup>1</sup>



Not all patients taking PDE5i are able to perform sexually<sup>1</sup>

## Clinical Data Confirms Role of VED\* in Combination with PDE5i<sup>2</sup>



**69 Men Aged 36 – 82**



**Max Dose of at Least 1 PDE5i**



**4 – 6 Attempts at Intercourse Over 3 Months**

### Baseline Statistics (PDE5i only)

- **Mean IIEF-5 Score = 9.0**<sup>\*\*</sup>
- **49% had erections inadequate for penetration**
- **72% had erections that did not last long enough for intercourse**
- **61% “not at all” or only “slightly” improved on PDE5i**

\*Vacuum Erection Device


\*\*International Index of Erectile Function

# Use of ErecAid<sup>®</sup> results in statistically significant improvements in all important erectile function dimensions<sup>2</sup>


## Initial Evaluation

## 4 Weeks Later with ErecAid


Were you able to insert your penis into your partner's vagina?



Did your erection last long enough to complete intercourse?



Has the treatment improved your erection?



**When PDE5i alone doesn't provide sufficient response use of VED provides for satisfactory intercourse.<sup>2</sup>**

Some patients should not use this device. Men who have a history of spontaneous priapism or blood disorders that can predispose them to priapism should not use the Osbon ErecAid. Men who have a bleeding disorder or bruise easily should also not use this device. Patients who are taking anticoagulants, have a penile implant, have limited manual dexterity, are unable to sense pain or pressure at the base of the penis, or have Peyronie's Disease should consult with their physician before using the ErecAid.



1. www.viagra.com. Accessibility verified November 3, 2011  
 2. Canguven O, Bailen J, Fredriksson W, Bock D, Burnett A. Combination of vacuum erection device and PDE5 inhibitors as salvage therapy in PDE5 inhibitor nonresponders with erectile dysfunction. J Sex Med. 2009;6:2561-2567

Contact your local Timm Medical Representative for Personalized Patient Training



9600 West 76th Street, Suite T  
 Eden Prairie, MN 55344 USA

800-438-8592  
 Phone: 952-947-9410  
 Fax: 952-947-9411  
 www.timmmedical.com