We are always happy to discuss any problems that you may have with the use of the Rejoyn System. If you have questions or would like to order replacement parts, please call our user help line 1-800-279-7434.

Any of the above listed treatments should be under a doctors care and/or by prescription.

- Vasedial Reconstructive Surger
  Prosthetic Implants (surgical)
- 5. Vascular Reconstructive Surgery
- 4. Intraurethral
- 3. Medications (oral and injectable)
- 2. Hormonal Therapy
- 1. Counseling

Loading Sleeve

auon

Other accepted forms of treatment for Impotence are:

NOTE: Carefully read instructions completely before use of the device.

### **BING KIL**

The support ring kit is used to maintain penile rigidity in men with erectile dysfunction by restricting penile venous outflow after the patient has obtained an erection (either with the aid of a vacuum pump or naturally). The support ring is placed around the base of the erect penis to restrict the flow of venous blood

leaving the penis.

The Ring Kit consists of:

Plus Zippered Pouch

Water Soluble Lubricant sgnift size Kings

WARNING: If you have symptoms of erectile dysfunction, (i.e., unable to get an erection that is good enough for sexual intercourse), talk to your doctor prior to using this device to avoid a possible harmful delay in figuring out any of the most common causes of this condition, such as diabetes, multiple sclerosis, cirrhosis of the liver, chronic renal failure, or alcoholism.

**WARNING:** Consult your doctor should any problems occur, and stop using the device if such conditions continue.

## Ring Kit

**POS-T-VAC MEDICAL** 

2111 W Wyatt Earp • PO Box 1436 Dodge City, KS 67801 Toll Free: 1-800-279-7434 Fax: 620-227-8474

## USER'S MANUAL

### Table of Contents

/	D
	iiX gniA shT seU oT woH
Е	Risks
7 ' I	Warnings & Precautions
ſ	Rejoyn Ring Kit Contents

Concerns

cause personal injury WARNING: If you do not heed the warnings below you may

- permanent injury to the penis. longer than 30 minutes at a time, the support ring may cause 1. Remove support ring within thirty (30) minutes. If worn
- .(səis base of the penis (where the shaft of the penis meets the pubic 2. Frequent use of support rings may result in bruising at the

# HOM TO USE YOUR RING KIT

.gniwerb loading sleeve, using your thumb nails as shown in press the support ring down over the cone onto the the entire surface of the cone. Urip each handle, and 1. Place a generous amount of water soluble lubricant on 6niA

Loading Sleeve

7

cause personal injury WARNING: If you do not heed these instructions you may

- permanent injury to the penis. longer than 30 minutes at a time, the support rings may cause Allow a minimum of 60 minutes between uses. If worn .1
- 2. Do not fall asleep wearing the support ring.
- Support rings should not be used as a birth control method. .ε
- Use the largest ring which holds an erection.
- .ς
- base of the penis (where the shaft of the penis meets the pubic 6. Frequent use of support rings may result in bruising at the Do not use support rings while using alcohol or drugs.
- Priapism (prolonged erection), and blocked urethra (urinary that causes distortion or abnormal bend of the erect penis), medical conditions such as Peyronie's disease ( a condition 7. Misuse of the support rings may worsen already existing .(səis
- . sinsq sht to 8. Do not use if you have decreased sensation of pain in the area tract).
- the removal of the rings difficult. 9. Do not use if you have decreased hand strength that will make
- rings and other latex type material such as condoms. 10.Do not use oil based lubricants they will be harmful to the
- soluble jelly when removing the support ring. red spots below the skin. Use caution and plenty of water concerned if you notice some bruising of the penis or small 11. Use of the support ring may cause bruising. Do not be overly

- 2. Separate the cone from the loading sleeve.
- your penis as close to your body as possible. the ring off the loading sleeve onto the base of 3. Slip the loading sleeve over the penis and slide
- comfortably. 4. The ring should grip the base of your penis
- .guri goes down, you may need to use a smaller 5. If you get a good firm erection and then it slowly
- bag is air free and tightly sealed for longer life. return the ring to its plastic bag. Make sure the side handles. Following use, wash, dry and 6. Remove the ring by pulling outward on the
- more information. Call our Toll Free number 1-800-279-7434 for may need to order the Rejoyn Vacuum System. vaginal entry when properly lubricated, you erection is not firm enough for satisfactory after following steps 1 through 5, and your 7. If you are unable to get a good, firm erection

Е

#### Don't Be Overly Concerned If...

disappear when you remove the support ring. increase of blood in the penis. The bluish color should the support ring. This may be a normal response to the Your penis turns slightly bluish in color after you apply

fpe support ring. is a normal response to the reduced blood flow caused by of the penis. This coolness is normal and not harmful. It You or your partner notice a slight drop in the temperature

for successful intercourse. find it necessary to try different positions with your partner its base when the ring is in place. If this occurs, you may Your erection has a "hinge" effect. The penis may pivot at

birth control and should not be used to avoid pregnancy. ring. Support rings are not a means of contraception or climax. Your semen may flow out when you remove the ejaculation process. This is not harmful, and does not affect The pressure of the support ring may interfere with the Your semen does not flow through at point of climax.